



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

# Journal Pre-proof

COVID-19 and non-traditional mask use: How do various materials compare in reducing the infection risk for mask wearers?

Amanda M. Wilson, Sarah E. Abney, Marco-Felipe King, Mark H. Weir, Martín López-García, Jonathan D. Sexton, Stephanie J. Dancer, Jessica Proctor, Catherine J. Noakes, Kelly A. Reynolds

PII: S0195-6701(20)30276-0

DOI: <https://doi.org/10.1016/j.jhin.2020.05.036>

Reference: YJHIN 6047

To appear in: *Journal of Hospital Infection*


Received Date: 24 April 2020

Accepted Date: 26 May 2020

Please cite this article as: RRH: COVID-19 mask infection risk comparison

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 The Healthcare Infection Society. Published by Elsevier Ltd. All rights reserved.



**COVID-19 and non-traditional mask use: How do various materials compare in reducing the infection risk for mask wearers?**

Running title: COVID-19 mask infection risk comparison

Amanda M. Wilson<sup>1\*</sup>, Sarah E. Abney<sup>1,2</sup>, Marco-Felipe King<sup>3</sup>, Mark H. Weir<sup>1,4</sup>, Martín López-García<sup>5</sup>, Jonathan D. Sexton<sup>1</sup>, Stephanie J. Dancer<sup>6,7</sup>, Jessica Proctor<sup>3</sup>, Catherine J. Noakes<sup>3</sup>, Kelly A. Reynolds<sup>1</sup>

1. Mel and Enid Zuckerman College of Public Health, University of Arizona, Tucson, AZ, USA
2. College of Agriculture and Life Sciences, University of Arizona, Tucson, AZ
3. School of Civil Engineering, University of Leeds, Woodhouse Lane, LS29JT, Leeds, UK
4. College of Public Health, The Ohio State University, Columbus, OH, USA
5. School of Mathematics, University of Leeds, Woodhouse Lane, LS29JT, Leeds, UK
6. School of Applied Sciences, Edinburgh Napier University, Edinburgh, UK
7. Department of Microbiology, Hairmyres Hospital, NHS Lanarkshire, G758RG, UK

\*Please address correspondence to Amanda M. Wilson, [apeifer@email.arizona.edu](mailto:apeifer@email.arizona.edu), 1295 N. Martin Ave. Drachman Hall PO Box 245210, Tucson, AZ

The SARS-CoV-2 pandemic has increased demands for surgical and respirator masks for healthcare workers (HCWs) and other frontline staff. The debate over the importance of airborne transmission of SARS-CoV-2 continues, but SARS-CoV-2 has been detected in air and laboratory studies demonstrate viability >12 hours in aerosols [1–3]. Low sampling volumes, air outlet fan location, and potential virus damage during sampling may explain SARS-CoV-2 detection variability [1,3].

Limited mask supply creates a risk for HCW exposure to SARS-CoV-2. Non-traditional materials are widely recommended for public use (source control) and have been considered in place of regulated masks in healthcare, especially in social-care settings. While various materials are effective for filtering large droplets, aerosols generated from sneezing, coughing, and aerosol-generating procedures may more readily pass through materials or leakage points [4]. There is a small amount of data on filtration efficacy but currently no quantitative modelling of efficacies for infection risk reduction.

We developed a probabilistic model to estimate infection risks for short (30-second, brief patient check) and long (20-minute, the duration required for patient intubation) inhalation exposure scenarios. These included situations in a COVID-19 patient room in which no mask was worn; an FFP2 (N95), FFP3 (N99) respirator or surgical mask was worn; or a non-traditional material mask (silk, tea towel, vacuum cleaner bag, pillowcase, antimicrobial pillowcase, cotton mix, 100% cotton t-shirt, linen, and scarf) was worn.

Inhaled viral dose was estimated using published concentrations (RNA/m<sup>3</sup>) of SARS-CoV-2 for >4 and 1-4 µm droplets measured in a hospital setting [1]. We used ranges from reported concentration data originating from a symptomatic and an asymptomatic patient to calculate minimum and maximum of randomly sampled uniform distributions [1]. Viral exposure for these two size ranges were summed to estimate a total inhaled dose. Doses were estimated for three assumed infectious fractions of total detected viral RNA: 0.1%, 1%, and 10%. Inhaled volumes (m<sup>3</sup>) were estimated using inhalation rates for men and women, where 5th and 99th percentiles of inhalation rates offered the uniform distribution minimum and maximum, respectively [5].

Filtration efficacies (fraction of total virus filtered out by the material) were used to model the reduction in viral inhalation exposure per material type. Due to lack of particle size-specific filtration efficacy data for these materials, we assumed filtration efficacy distributions were applicable to both particle size ranges. For each 10,000 combinations investigated, a filtration efficacy was randomly sampled from a normal distribution, left- and right-truncated at 0 and 1, respectively. For surgical mask and non-traditional materials, mean and standard deviations of efficacies were informed by MS2 filtration efficacies [6]. Mean efficacies of 95% and 99% were assumed for FFP2 and FFP3 respirators, respectively. SDs were provided by Rengasamy et al. (2009), where larger SDs of two manufacturer versions were chosen as a conservative risk approach [7].

Data from SARS-CoV-1 and human coronavirus 229E (HCoV-229E) dose-response curves were used to estimate a SARS-CoV-2 exact beta-Poisson curve [8]. Based on current epidemiological knowledge, we assume the infectivity of SARS-CoV-2 lies between SARS-CoV-

1 and HCoV-229E. Pairs of bootstrapped  $\alpha$  and  $\beta$  values were used to estimate infection risk per dose.

Comparing no protection (baseline) for 20-minute and 30-second exposures, we predicted that mean infection risks were reduced by 24 - 94% and by 44 - 99% depending on the mask. Risk reductions decreased as exposure durations increased. The greatest reduction in estimated mean infection risk was for FFP3 masks, which as expected reduced baseline mean risks by 94% and 99% for 20-minute and 30-second exposures, respectively (Figure 1). Of non-traditional materials, the vacuum cleaner bag resulted in the greatest reduction in mean infection risk (20-minute exposure: 58%, 30-second exposure: 83%), while scarves offered the lowest (20-minute exposure: 24%, 30-second exposure: 44%) (Figure 1). However, large filtration variability, such as for silk or the tea towel, should be considered when comparing non-traditional mask materials (Figure 1).

Limitations include not accounting for viral transfer from the hands to the mask during mask adjustments and assuming all masks were worn in the same way. Realistically, homemade mask fit is likely to be more variable than for regulated masks. While the HCoV-229E data utilized for the dose-response curve was based on human data, the SARS-CoV-1 dose-response data originated from an animal-feeding study [8]. Future work includes updating the dose-response curve as data on SARS-COV-2 emerges and addressing the effects of design/fit on infection risk.

We demonstrate that some materials, like vacuum cleaner bags, may be effective alternatives for reducing infection risk. While N95 masks (and similar respirators) are recommended for HCWs and others in close proximity to aerosol generating procedures, alternative materials may be useful where there are PPE shortages. This may be of particular relevance in low resource settings where access to PPE is considerably more limited.

### **Code, Materials**

Under a Creative Commons Zero v1.0 Universal license (CC-BY), code can be accessed at: <https://github.com/awilson12/COVID-19-Mask-Note>

### **Conflicts of Interest**

The authors have no conflicts of interest to declare.


### **Funding Source**

A.M. Wilson was supported by the University of Arizona Foundation and the Hispanic Women's Corporation/Zuckerman Family Foundation Student Scholarship Award through the Mel and Enid Zuckerman College of Public Health, University of Arizona. M-F. King and C.J. Noakes were funded by the Engineering and Physical Sciences Research Council, UK: Healthcare Environment Control, Optimisation and Infection Risk Assessment (<https://HECOIRA.leeds.ac.uk>) (grant code: EP/P023312/1). M. López-García was funded by the Medical Research Council, UK (MR/N014855/1). J. Proctor was funded by EPSRC Centre for Doctoral Training in Fluid Dynamics at Leeds (grant code EP/L01615X/1). S.E. Abney was

funded by a research assistantship from the US-Israel Binational Agricultural Research Development Fund and through a University of Arizona Graduate Access Scholarship.

## References

- [1] Chia P, Coleman K, Tan Y, Ong S, Gum M, Lau S, et al. Detection of air and surface contamination by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) in hospital rooms of infected patients 2020. Preprint. <https://doi.org/https://doi.org/10.1101/2020.03.29.20046557>.
- [2] Fears A, Klimstra W, Duprex P, Hartman A, Weaver S, Plante K, et al. Comparative dynamic aerosol efficiencies of three emergent coronaviruses and the unusual persistence of SARS-CoV-2 in aerosol suspensions 2020. Preprint. <https://doi.org/https://doi.org/10.1101/2020.04.13.20063784>.
- [3] Ong SWX, Tan YK, Chia PY, Lee TH, Ng OT, Wong MSY, et al. Air, surface environmental, and personal protective equipment contamination by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) from a symptomatic patient. *JAMA* 2020;2–4. <https://doi.org/10.1001/jama.2020.3227>.
- [4] Weber A, Willeke K, Marchloni R, Myojo T, McKay R, Donnelly J, et al. Aerosol penetration and leakage characteristics of masks used in the health care industry. *Am J Infect Control* 1993;21:167–73. [https://doi.org/10.1016/0196-6553\(93\)90027-2](https://doi.org/10.1016/0196-6553(93)90027-2).
- [5] U.S. Environmental Protection Agency. Exposure Factors Handbook 2011 Edition (EPA/600/R-09/052F). Washington, DC: 2011.
- [6] Davies A, Thompson KA, Giri K, Kafatos G, Walker J, Bennett A. Testing the efficacy of homemade masks: would they protect in an influenza pandemic? *Disaster Med Public Health Prep* 2013;7:413–8. <https://doi.org/10.1017/dmp.2013.43>.
- [7] Rengasamy S, Eimer BC, Shaffer RE. Comparison of nanoparticle filtration performance of NIOSH-approved and CE-marked particulate filtering facepiece respirators. *Ann Occup Hyg* 2009;53:117–28. <https://doi.org/10.1093/annhyg/men086>.
- [8] Watanabe T, Bartrand TA, Weir MH, Omura T, Haas CN. Development of a dose-response model for SARS coronavirus. *Risk Anal* 2010;30:1129–38. <https://doi.org/10.1111/j.1539-6924.2010.01427.x>.



**Figure 1.** Distributions of estimated infection risks for FFP3, FFP2, surgical masks, masks made of non-traditional materials (vacuum cleaner bag, tea towel, cotton mix, antimicrobial pillowcase, linen, pillowcase, silk, 100% cotton T-shirt, scarf) and no mask for 30 seconds (0.5 minutes) or 20 minutes of inhalation exposure\*

\*Vertical lines indicate the 25th, 50th, and 75th infection risk percentiles